


#### **Application Instructions**







2. Apply skin prep evenly to areas adhesive will adhere to, and let dry for approximately 30 seconds.


1. Secure and dress IV line per hospital protocol.

- 3. Center the pad over the IV & dressing. Determine if length of legs need to be cut to prevent overlapping. If necessary, cut legs before removing liner.
  - \*Do not cut more than 2" from one side. If you require cutting more than 2", you must trim from other set of product legs.




4. Pull back liner and press product to adhere to patient in one motion, one side at a time. Do not stretch to apply.

## **Antecubital Application**



### **Hand Application**



### **Forearm Application**



To inspect IV insertion site, lift center section of product to visualize area.

#### ORDERING INFORMATION **Product Number** Size Quantity 675 One size fits most 10/Box

HCPCS Code A4221

# **Product Features**

- Helps maintain a continuous IV flow while protecting and preserving the IV site
- Reduces the number of IV restarts, which decreases risk of infection
- Saves nursing time and reduces alarm fatigue
- Allows for patient movement while reducing downstream occlusion alarms caused by movement
- Improves patient comfort without the need for a stiff immobilizing device
- Reduces risk of dislodgement from picking and pulling of the IV by confused patients

Keep device clean and dry. Change after 3 days or sooner if necessary.

Removal: Device should be removed with care to avoid skin trauma. Re-apply skin prep if another device is to be applied.

